

(ii) A donor with a reactive serologic test for syphilis shall not be plasmapheresed again until the donor's serum is tested and found to be non-reactive to a serologic test for syphilis, except as provided in paragraph (b)(2) (iii) and (iv) of this section.

(iii) A donor whose serum is determined to have a biologic false-positive reaction to a serologic test for syphilis may be plasmapheresed: *Provided*, That the donor's file identifies the serologic test for syphilis and results used to confirm the biologic false-positive reaction and indicates that the physician on the premises has determined the false-positive reaction is not the result of an underlying disorder that would disqualify the donor from participation in the plasmapheresis program. If the serologic test for syphilis is performed at a facility other than the plasmapheresis center, all applicable provisions of § 640.71 shall be met.

(iv) A donor with a reactive serologic test for syphilis may be plasmapheresed only to obtain plasma to be used for further manufacturing into control serum for the serologic test for syphilis: *Provided*, That the physician on the premises approves the donation, the donor's file contains a signed statement from a physician or clinic establishing that treatment for syphilis has been initiated and that continuance in the plasmapheresis program will not interfere with or jeopardize the treatment of the syphilitic donor.

(3) A donor identification system shall be established that positively identifies each donor and relates such donor directly to his blood and its components as well as to his accumulated records and laboratory data. Such system shall include either a photograph of each donor which shall be used on each visit to confirm the donor's identity, or some other method that provides equal or greater assurance of positively identifying the donor.

(4) The amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure or in any 2-day period shall not exceed 1,000 milliliters unless the donor's weight is 175 pounds or greater, in which case the amount of whole blood, not including anticoagu-

lant, removed from the donor during a manual plasmapheresis procedure or in any 2-day period shall not exceed 1,200 milliliters.

(5) The amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure within a 7-day period shall not exceed 2,000 milliliters unless the donor's weight is 175 pounds or greater, in which case the amount of whole blood, not including anticoagulant, removed from a donor during a manual plasmapheresis procedure within a 7-day period shall not exceed 2,400 milliliters.

(6) No more than 500 milliliters of whole blood shall be removed from a donor at one time, unless the donor's weight is 175 pounds or greater, in which case no more than 600 milliliters of whole blood shall be removed from the donor at one time.

(7) The plasma shall be separated from the red blood cells immediately after blood collection. The maximum feasible volume of red blood cells shall be returned to the donor before another unit is collected.

(8) The volume of plasma collected during an automated plasmapheresis collection procedure shall be consistent with the volumes specifically approved by the Director, Center for Biologics Evaluation and Research, and collection shall not occur less than 2 days apart or more frequently than twice in a 7-day period.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 10769, Mar. 12, 1976; 64 FR 45373, Aug. 19, 1999; 64 FR 56453, Oct. 20, 1999]

§ 640.66 Immunization of donors.

If specific immunization of a donor is to be performed, the selection and scheduling of the injection of the antigen, and the evaluation of each donor's clinical response, shall be by a qualified licensed physician or physicians. The administration of the antigen may be performed by a licensed physician or a trained person under his supervision. Any material used for immunization shall be either a product licensed under section 351 of the Public Health Service Act for such purpose or one specifically approved by the Director, Center for Biologics Evaluation and Research,

§ 640.67

Food and Drug Administration. Immunization procedures shall be on file at each plasmapheresis center where immunizations are performed.

[38 FR 32089, Nov. 20, 1973, as amended at 49 FR 23834, June 8, 1984; 55 FR 11013, Mar. 26, 1990]

§ 640.67 Laboratory tests.

Each unit of Source Plasma shall be tested for evidence of infection due to communicable disease agents as required under § 610.40 of this chapter.

[66 FR 31165, June 11, 2001]

§ 640.68 Processing.

(a) *Sterile system.* All administration and transfer sets inserted into blood containers used for processing Source Plasma intended for manufacturing into injectable or noninjectable products and all interior surfaces of plasma containers used for processing Source Plasma intended for manufacturing into injectable products shall be sterile, pyrogen-free, nontoxic, and compatible with the contents under normal conditions of use. Only Sodium Chloride Injection USP shall be used as a red blood cell diluent. If the method of separation of the plasma intended for injectable products involves a system in which an airway must be inserted into the plasma container, the airway shall be sterile and constructed so as to exclude microorganisms and maintain a sterile system.

(b) *Final containers.* Final containers used for Source Plasma, whether integrally attached or separated from the original blood container, shall not be entered prior to issuance for any purpose except for filling with the plasma. Such containers shall be uncolored and hermetically sealed, and shall permit clear visibility of the contents. Final containers and their components shall not interact with the plasma contents under conditions of storage and use so as to alter the safety, quality, purity, or potency of the plasma and shall provide adequate protection against external factors that may cause deterioration or contamination. Prior to filling, the final container shall be marked or identified by number or other symbol which will relate it directly to the donor.

21 CFR Ch. I (4–1–02 Edition)

(c) *Preservative.* Source Plasma shall not contain a preservative.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 10769, Mar. 12, 1976; 50 FR 4140, Jan. 29, 1985]

§ 640.69 General requirements.

(a) *Pooling.* Two units of Source Plasma from the same donor may be pooled if such units are collected during one plasmapheresis procedure: *Provided*, That the pooling is done by a procedure that does not introduce a risk of contamination of the red blood cells and, for plasma intended for injectable products, gives maximum assurance of a sterile container of plasma.

(1) The pooling of plasma from two or more donors is not permitted in the manufacture of Source Plasma intended for manufacturing into injectable products.

(2) The pooling of plasma from two or more donors by the manufacturer of Source Plasma intended for manufacturing into noninjectable products is permitted: *Provided*, That the plasma from two or more donors is pooled after the plasma has been removed from the red blood cells, and after the red blood cell containers are sealed.

(b) *Storage.* Immediately after filling, plasma intended for manufacturing into injectable products shall be stored at a temperature not warmer than -20°C , except for plasma collected as provided in § 640.74. Plasma intended for manufacturing into noninjectable products may be stored at temperatures appropriate for the intended use of the final product, provided these temperatures are included in the Source Plasma license application.

(c) *Inspection.* Source Plasma intended for manufacturing into injectable products shall be inspected for evidence of thawing at the time of issuance, except that inspection of individual plasma containers need not be made if the records of continuous monitoring of the storage temperature establish that the temperature remained at -20°C or colder. If there is evidence that the storage temperature has not been maintained at -20°C or colder, the plasma may be relabeled and issued as provided in § 640.76(a).